

Amendment to the Specification

Please replace the paragraph starting on page 29, line 17, through page 30, line 9, of the specification with the following rewritten paragraph:

The fifth column of Table 1B.1, "ORF (From-To)", provides the location (i.e., nucleotide position numbers) within the polynucleotide sequence of SEQ ID NO:X that delineates the preferred open reading frame (ORF) that encodes the amino acid sequence shown in the sequence listing and referenced in Table 1B.1 as SEQ ID NO:Y (column 6). Column 7 of Table 1B.1 lists residues comprising predicted epitopes contained in the polypeptides encoded by each of the preferred ORFs (SEQ ID NO:Y). Identification of potential immunogenic regions was performed according to the method of Jameson and Wolf (CABIOS, 4; 181-186 (1988)); specifically, the Genetics Computer Group (GCG) implementation of this algorithm, embodied in the program PEPTIDESTRUCTURE (Wisconsin Package v10.0, Genetics Computer Group (GCG), Madison, Wisc.). This method returns a measure of the probability that a given residue is found on the surface of the protein. Regions where the antigenic index score is greater than 0.9 over at least 6 amino acids are indicated in Table 1B.1 as "Predicted Epitopes". In particular embodiments, polypeptides of the invention comprise, or alternatively consist of, one, two, three, four, five or more of the predicted epitopes described in Table 1B.1. It will be appreciated that depending on the analytical criteria used to predict antigenic determinants, the exact address of the determinant may vary slightly. Column 8 of Table 1B.1 ("Tissue Distribution") is described below in Table 1B.2 Column 5. Column 9 of Table 1B.1 ("Cytologic Band") provides the chromosomal location of polynucleotides corresponding to SEQ ID NO:X. Chromosomal location was determined by finding exact matches to EST and cDNA sequences contained in the NCBI (National Center for Biotechnology Information) UniGene database. Given a presumptive chromosomal location, disease locus association was determined by comparison with the Morbid Map, derived from Online Mendelian Inheritance in Man (Online Mendelian Inheritance in Man, OMIM™, McKusick-Nathans Institute for Genetic Medicine, Johns Hopkins University (Baltimore, MD) and National Center for Biotechnology Information, National Library of Medicine (Bethesda, MD) 2000. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>). If the putative chromosomal location of the Query overlaps with the chromosomal location of a Morbid Map entry, an OMIM identification number is disclosed in Table 1B.1, column 9 labeled "OMIM Disease Reference(s)". A key to the OMIM reference identification numbers is provided in Table 5.

Please replace the paragraph on page 35, lines 17-20 of the specification with the following rewritten paragraph:

The "Exemplary Accessions" Column indicates GenBank Accessions (available online through the National Center for Biotechnology Information (NCBI) at

<http://www.ncbi.nlm.nih.gov/>) which correspond to the "Exemplary Targets" shown in the adjacent row.

Please replace the paragraph starting on page 37, line 34, through page 38, line 3 of the specification with the following rewritten paragraph:

Table 5 provides a key to the OMIM reference identification numbers disclosed in Table 1B.1, column 9. OMIM reference identification numbers (Column 1) were derived from Online Mendelian Inheritance in Man (Online Mendelian Inheritance in Man, OMIM. McKusick-Nathans Institute for Genetic Medicine, Johns Hopkins University (Baltimore, MD) and National Center for Biotechnology Information, National Library of Medicine, (Bethesda, MD) 2000. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>). Column 2 provides diseases associated with the cytologic band disclosed in Table 1B.1, column 8, as determined using the Morbid Map database.

Please replace the paragraph on page 3438, lines 15-18 of the specification with the following rewritten paragraph:

The "Exemplary Accessions" Column indicates GenBank Accessions (available online through the National Center for Biotechnology Information (NCBI) at <http://www.ncbi.nlm.nih.gov/>) which correspond to the "Exemplary Targets" shown in the adjacent row.

Please replace the paragraph on page 3947, lines 4-10 of the specification with the following rewritten paragraph:

Table 5 provides a key to the OMIM reference identification numbers disclosed in Table 1B.1, column 9. OMIM reference identification numbers (Column 1) were derived from Online Mendelian Inheritance in Man (Online Mendelian Inheritance in Man, OMIM. McKusick-Nathans Institute for Genetic Medicine, Johns Hopkins University (Baltimore, MD) and National Center for Biotechnology Information, National Library of Medicine, (Bethesda, MD) 2000. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>). Column 2 provides diseases associated with the cytologic band disclosed in Table 1B.1, column 8, as determined using the Morbid Map database.

Please replace the paragraph starting on page 3981, line 33, through page 3982, line 5 of the specification with the following rewritten paragraph:

Any polypeptide sequence encoded by, for example, the polynucleotide sequences set forth as SEQ ID NO:X or the complement thereof, (presented, for example, in Tables 1A and 2), the cDNA

contained in ATCC Deposit No:Z, or the polynucleotide sequence as defined in column 6 of Table 1C, may be analyzed to determine certain preferred regions of the polypeptide. For example, the amino acid sequence of a polypeptide encoded by a polynucleotide sequence of SEQ ID NO:X (e.g., the polypeptide of SEQ ID NO:Y and the polypeptide encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2) or the cDNA contained in ATCC Deposit No:Z may be analyzed using the default parameters of the DNASTAR computer algorithm (DNASTAR, Inc., 1228 S. Park St., Madison, WI 53715 USA; <http://www.dnastar.com/>).

Please replace the cell in Table 1D on page 3410 of the specification with the following rewritten cell:

614	HWHGU54	1562	<p>Production of IL-8 by by-endothelial cells (such as Human Umbilical Cord Endothelial Cells).</p>	<p>Assays measuring production of IL-8 are well known in the art and may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate production and/or secretion of IL-8. For example, FMA7 may be used or routinely modified to assess the ability of polypeptides of the invention (including antibodies and agonists or antagonists of the invention) to regulate production and/or secretion of IL-8 from endothelial cells (such as human umbilical vein endothelial cells (HUVEC)). HUVECs are endothelial cells which line venous blood vessels, and are involved in functions that include, but are not limited to, angiogenesis, vascular permeability, vascular tone, and immune cell extravasation. Endothelial cells play a pivotal role in the initiation and perpetuation of inflammation and secretion of IL-8 may play an important role in recruitment and activation of immune cells such as neutrophils, macrophages, and lymphocytes.</p>	<p>Highly preferred indications include immunological and inflammatory disorders (e.g., such as allergy, asthma, leukemia, etc. and as described below under "Immune Activity", and "Blood-Related Disorders"). Highly preferred indications also include autoimmune disorders (e.g., rheumatoid arthritis, systemic lupus erythematosus, Crohn's disease, multiple sclerosis and/or as described below), neoplastic disorders (e.g., organ cancers such as lung, liver, colon cancer, and/or as described below under "Hyperproliferative Disorders"), and cardiovascular disorders (e.g., such as described below under "Cardiovascular Disorders"). Preferred indications include thrombosis, bacteremia and sepsis syndrome and consequent complications (such as acute respiratory distress syndrome and systemic ischemia-reperfusion resulting from septic shock), restenosis and atherosclerosis.</p>
-----	---------	------	---	--	---